



TITLE:

Fragmentation of multiply ionized biomolecules induced by MeV heavy ions

AUTHOR(S):

Yoshida, S. O.; Majima, T.; Asai, T.; Matsubara, M.; Tsuchida, H.; Itoh, A.

CITATION:

Yoshida, S. O. ...[et al]. Fragmentation of multiply ionized biomolecules induced by MeV heavy ions. Journal of Physics: Conference Series 2015, 635: 032062.

ISSUE DATE:

2015-09-07

URL:

<http://hdl.handle.net/2433/223216>

RIGHT:

© Published under licence by IOP Publishing Ltd.; Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Fragmentation of multiply ionized biomolecules induced by MeV heavy ions

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2015 J. Phys.: Conf. Ser. 635 032062

(<http://iopscience.iop.org/1742-6596/635/3/032062>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 130.54.110.31

This content was downloaded on 27/04/2017 at 02:45

Please note that [terms and conditions apply](#).

You may also be interested in:

[Electron impact cross-sections for biomolecules - completeness and self-consistency via swarm analysis](#)

J de Urquijo, M Casey, D Konvalov et al.

[Electron and positron scattering from biomolecules](#)

M J Brunger, S J Buckman and A Zecca

[Preparation of 1D nanostructures using biomolecules](#)

Stela Pruneanu, Liliana Olenic, Lucian Barbu Tudoran et al.

[Ultrafast nonadiabatic fragmentation dynamics of biomolecules](#)

Pablo López-Tarifa, Dariusz Grzegorz, Piekarski et al.

[Structure and Function of Biomolecules](#)

Marek Cieplak and Andrzej Sienkiewicz

[Synthesis of new pyrrole-containing biomolecules as building blocks for functionalized polypyrroles in nanobiotechnology](#)

S Karsten, A Nan, R Turcu et al.

[Microcantilever-Based Biosensor for Detection of Various Biomolecules](#)

Kyung-Ah Yoo, Kwang-Ho Na, Seung-Ryong Joung et al.

[A new apparatus for the study of electron impact fragmentation of molecular clusters](#)

G Barrett and P J M van der Burgt

[Molecular and Nanodynamics: from atoms to biomolecules](#)

Vincenzo Aquilanti

Fragmentation of multiply ionized biomolecules induced by MeV heavy ions

S. O. Yoshida*, T. Majima^{*†1}, T. Asai*, M. Matsubara*, H. Tsuchida^{*†} and A. Itoh^{*†}


* Department of Nuclear Engineering, Kyoto University, Kyoto 615-8540, Japan

† Quantum Science and Engineering Center, Kyoto University, Uji 611-0011, Japan

Synopsis Collision-induced ionization and fragmentation of gas-phase adenine molecules ($C_5N_5H_5$) have been investigated for single electron capture collisions of 1.2 MeV C^{2+} ions. Coincidence measurements between the number of ejected electrons and the time-of-flight of fragment ions allows us to determine charge states of intermediate parent ions r . Highly charged parent ions are observed up to $r \sim 6$ and the production of lighter fragment ions becomes predominant with increasing r . We found that kinetic energies of H^+ are significantly larger than those of other fragment ions and become nearly equivalent irrespective of r for $r > 1$.

Ion-induced fragmentation of DNA building blocks has gained increasing attention as fundamental processes in particle cancer therapy and radiation damage. In MeV heavy ion collisions, a large amount of energies are transferred from an incident ion to target electrons, leading to highly electronic excitation and multiple ionization of the target molecule. To understand the details of molecular fragmentation, we have extensively studied so far on the correlation between charged states of intermediate parent ions and fragmentation patterns for various polyatomic molecules [1, 2]. In this work, we extend measurements to an isolated biomolecular target of adenine ($C_5N_5H_5$).

The experimental procedure is given elsewhere [1]. Correlation data between charge states of intermediate parent ions and kinetic energies (KE) of fragment ions have been obtained under a single electron capture condition of 1.2 MeV C^{2+} ions. The reaction is written as



where r represents the charge state of intermediate parent ions and is determined directly from the number of ejected electrons n_e : $r = n_e + 1$.

Figure 1 shows a result of total time-of-flight (TOF) mass spectrum of fragment ions. The main peaks are H^+ and CNH_2^+ . Compared with slow ion collisions [3], relative yields of smaller fragment ions are high because of a large amount of energy transfers. It should be pointed out that fragment ions such as CNH_3^+ , $CN_2H_3^+$ and $C_2N_3H_4^+$ are produced from proton migration in a molecule. Multiple ionization of up to $r \sim 6$ was observed by a direct electron counting technique [1-3]. By analyzing the coincidence data between the TOF and the number of ejected electrons, partial TOF spectra associated with a specific value of r were obtained. It is found that the intensities of lighter fragment ions increase with increasing r .




Figure 1. Total TOF spectrum of fragment ions obtained in collisions of 1.2 MeV C^{2+} with isolated adenine molecules.

Most of the fragment ions gain KE less than 10 eV, while H^+ gains significantly large KE extending to several tens of eV. It indicates a possibility of further fragmentation of DNA by these “high energy” protons [4]. The KE distribution of H^+ varies greatly for different values of r . As for $r = 1$, it peaks at 0 eV, while for $r > 1$ the KE peaks at about 6 eV with a tail extending to about 40 eV. It is pointed out that KE distributions are nearly independent of r for $r > 1$. While the reason is not clear at present, the simple Coulomb explosion model may not hold in the present collision system.

The work was supported by JSPS KAKENHI Grants No.24340096, No. 23760826 and Mizuho Foundation for the Promotion of Sciences.

References

- [1] T. Majima *et al* 2014 *Phys. Rev. A* **90** 062711
- [2] T. Majima *et al* 2004 *Phys. Rev. A* **69**, 031202(R)
- [3] R. Brédy *et al* 2007 *Nucl. Instr. and Meth. B.* **261** 114
- [4] Z. Deng *et al* 2006 *Phys. Rev. Lett.* **96** 243203

¹ E-mail: majima@nucleng.kyoto-u.ac.jp

